arXiv:1603.04641v2 [cs.GT] 17 Jan 2017

COMPOSITIONAL GAME THEORY

NEIL GHANI, JULIAN HEDGES, VIKTOR WINSCHEL, AND PHILIPP ZAHN

University of Strathclyde
e-mail address: ng@cis.strath.ac.uk

University of Oxford
e-mail address: julian.hedges@cs.ox.ac.uk

OICOS GmbH
e-mail address: viktor.winschel@googlemail.com

University of St Gallen
e-mail address: philipp.zahn@unisg.ch

Abstract. This paper aims to significantly impact the way we conceive of, reason about, and construct software for economic game theory. We achieve this by building an new and original model of economic games based upon the computer science idea of compositionality: concretely we i) give a number of operators for building up complex and irregular games from smaller and simpler games; and ii) show how the equilibria of these complex games can be defined recursively from the equilibria of their simpler components. We apply compositional reasoning to sophisticated games where agents must reason about how their actions effect future games and how those future games effect the utility they receive. This requires further originality in that we augment the usual lexicon of games with a new concept of coutility. This notion — dual to utility — arises because, in order for games to accept utility, this utility must be generated by other games. A third source of originality is the representation of our games as string diagrams so as to give a clear visual interface to manipulate them. Our fourth, and final, source of originality is a categorical formalisation of these intuitive string diagrams which guarantees our model is mathematically rigourous.

Introduction

We must all grapple with this important question:

> *How can we ensure that our theories of the world scale from the small examples in our classrooms or on our blackboards, to the huge and complex systems that appear in the real world?*

Key words and phrases: Game theory, Economic equilibria, String diagrams, Monoidal categories, Selection functions.

LOGICAL METHODS
IN COMPUTER SCIENCE
DOI:10.2168/LMCS-???
© Ghani, Hedges, Winschel, and Zahn
Creative Commons

One answer — the brute force answer — is to simply be very, very good at applying our theories. For example in computer science, many researchers are grappling with the phenomena of big data by building faster and faster computers. The equivalent within economic game theory might be to represent complex games as huge pay-off matrices of actions and associated utilities and use some of those very fast computers to compute the associated Nash equilibria of these games. An alternative to brute force techniques, widely regarded within computer science as being best practice, is *compositionality* where one sees complex systems as being built from smaller subsystems. Because these subsystems are smaller, they are easier to reason about. One then combines properties of these subsystems into properties of the overall system. Compositionality also promotes reuse: if a particular system is a subsystem of many other systems, then any results concerning that system do not need to be regenerated whenever compositional reasoning takes place on any super-system of that system: this further promotes efficient reasoning. Examples of compositional reasoning are numerous, e.g.

- Within logic, the truth of a formulae $A \wedge B$ is computed from the truth of A and the truth of B which, in turn, are computed recursively.
- Within programming languages, the effect of executing a programme p *then* q is computed by combining the effect of executing p with the effect of executing q. These, in turn, are computed recursively.
- Within concurrency, Milner's highly influential *Communicating Sequential Processes* proposes operators for building processes. As above, properties of processes are proved by combining properties of sub-processes.

Can the gains of compositionality within Computer Science be replicated within economic game theory? After all, not all reasoning can be put in compositional form, especially if there is significant emergent behaviour present in a large system which is not present in subsystems. And, such emergent behaviour is certainly present within economic games. For example, if σ is an optimal strategy for a game G, then is σ part of an optimal strategy for $G * H$ where H is another game and $G * H$ is some super-game built from G and H? Clearly the answer is not necessarily! More concretely, the iterated prisoner's dilemma has many equilibria (such as cooperative equilibria) that do not arise from repeatedly playing the Nash equilibrium of the stage game. Indeed, the difficulty in constructing a compositional model of economic game theory led one economist present at a recent talk on this work to exclaim "*This can't be done!*" while another described this as "*a conjuring trick*" in that it seemed to achieve the impossible!

So, how do we achieve the impossible? Our approach to cutting this particular Gordian knot is to observe that, at its essence, any compositional model of game theory must model not just each game, but the interaction of each game with all other games it might interact with. Of course, all current models of economic games do not attempt this endeavour with good reason — as just demonstrated, this interaction is non-trivial and a priori, there is a huge number of such games - indeed a proper class! Our solution — and the crucial technical insight underpinning our model — is that it suffices to replace the class of games which a given game might interact with the utility functions of those games. Put simply, standard models of game theory hardwire a specific utility function into the definition of a game while the novelty of our model is that each game is parameterised by the space of *all possible utility* functions. This is accomplished by embedding within each game "ports" where interaction with other games takes place in the form of sending the environment information about moves played and receiving utility from the environment. This leads to a second innovation

within this paper: we introduce a new representation of games which complements the usual representation of games via trees or pay-off matrices. This new representation of games uses string diagrams because they enable us to clearly visualise the structure of complex games via the places where the ports of one game are glued together with the ports of other games. For example, a bimatrix game such as the Prisoner's Dilemma can be represented as the following string diagram



where i) X is the type of the current state of the world; ii) Y is the type of possible actions or moves; iii) R is the type of utilities or payoffs; and iv) S is the type of the coutility extruded back to the environment. This concept of *coutility* is another of our contributions — coutility arises as, in a closed system, the utility of one player must have been generated by another player. For example, if I place a bet with a bookmaker and expect utility in the form of winnings, those winnings will have to come from the bookmaker. This we call the bookmakers coutility. Pregames also possess functions describing how moves are chosen, exactly how much coutility to extrude, and a relation describing its equilibria. Crucially, the equilibria relation can be *any relation* and need not be restricted to Nash equilibria. The scientific value of these pregames is that they form a compositional model of economic games in that we can provide operators for building complex pregames from simpler pregames and, in particular, for defining the equilibria of these complex pregames from the equilibria of their component pregames. Our final contribution is a categorical interpretation of pregames and their string diagram representations which ensures they have precise mathematical meanings and thus that our reasoning about them is mathematically rigorous.

Related Work: The context of this paper begins with [15], which for the first time approached game theory using ideas from program semantics. Nevertheless, the game theory developed in that paper is no more compositional than ordinary game theory. A key point of originality within our approach is the use of continuations to allow players to consider the possible outcomes of decisions and factor those possible outcomes into their decision making process. Continuations are implicit in the literature on selection functions (see [7] for an introduction). Pregames are also closely related to the 'partially defined games' of [13]. Other significantly influential work on using logical methods in economics are [1, 11].

The relationship between string diagrams and monoidal categories has been developed extensively in quantum information theory [3] and bialgebra [5], and also applied to distributional semantics in linguistics [6]. There are many variants of monoidal categories, each with its own associated string language, surveyed in [18]. The language used in this paper is a fragment of the one for compact closed categories (section 4.8 of *loc. cit.*). We use

a fragment because, within game theory, there is only a limited form of backward-causality (arising from rational agents reasoning about potential future values).

The paper is structured as follows: Section 1 introduces some of the notation used in this paper while Section 2 demonstrates the feasibility of a compositional model of game theory by considering a simple class on non-dynamic games. Section 3 introduces pregames and their representation as string diagrams, Section 4 defines the key operators of parallel and sequential composition of pregames, Section 5 gives a rigorous mathematical foundation for pregames and their representation as string diagrams. Finally, Section 6 discusses the choices inherent within our compositional model and directions for further research.

1. Preliminaries

We denoted by $\mathbb{R}$ the set of real numbers. We also denote by 1 the set containing only the element $*$ - when a function requires an input of type 1, we often elide that input to increase legibility. We use the λ-calculus notation $\lambda x.t$ to denote the function which takes x as input and returns the value of t — this will of course usually depend upon x. If f is a function in two variables, we often write $f(-, x)$ and $f(x, -)$ respectively for the function in one variable obtained by instantiating the first and second inputs respectively of f to be x. We also follow computer science notation and write $f\ x$ — as opposed to $f(x)$ — for the application of a function f to an input x. Composition of functions is written $f \circ g$ and projections are denoted $\pi_1 : A \times B \to A$ and $\pi_2 : A \times B \to B$. We denote by $\mathcal{P}$ the powerset functor on sets. At the centre of game theory is the desire to maximise utility and so we define, for any function $f : A \to B$ (where B is a preordered set), the set $\mathsf{argmax} f$ of inputs where f attains a maximal value.

$$\mathsf{argmax}\ f = \{a \in A \mid \forall a' \in A. fa \geq fa'\}$$

We assume familiarity with basic category theory as can be found in [12]. Readers not familiar with these topics can simply skip the technical details of Section 6 — the paper has been written to be accessible to economists as well as computer scientists and hence the technical details have been so isolated.

2. Simple Games, Compositionality and Nash Equilibria

To ensure accessibility of our methods to non-computer scientists, we begin by treating simple games where players choose actions and utility is then generated. Without loss of generality, we consider just two players

Definition 2.1 (Simple 2-player Game)**.** A simple two-player game G consists of a 5-tuple $(Y_1, Y_2, R_1, R_2, k : Y_1 \times Y_2 \to R_1 \times R_2)$, where Y_i are the sets of moves available to each player, R_i are the types of utility expected by each players, and k computes the utility associated to each pair of moves. Let $k_1 = \pi_1 \circ f$ and $k_2 = \pi_2 \circ f$. Then, the set of *Nash equilibria* $E_G \subseteq Y_1 \times Y_2$ is defined by

$$\begin{aligned}(y_1, y_2) \in E \quad \mathsf{iff} \quad & y_1 \in \mathsf{argmax}\ k_1(-, y_2) \\ & \wedge\ y_2 \in \mathsf{argmax}\ k_2(y_1, -)\end{aligned}$$

This definition of Nash equilibrium is not compositional: it is not derived from more primitive concepts but rather is postulated as being itself a primitive concept whose myriad of different applications, and conceptual simplicity, underpin its value. While of course accepting its usefulness, and conceptual simplicity, we do believe it can be derived in a compositional manner from the interaction of the first player with the second player. Substantiating this belief will show that a compositional model of economic game theory is possible and what it might look like. First, consider the players:

Definition 2.2 (Simple Game)**.** A simple game consists of a set Y of moves, a set R of utilities and an equilibria function $E : (Y \to R) \to \mathcal{P}Y$. The set of simple games with actions Y and utilities R is written $\mathsf{SP}_R Y$

In the above definition $\mathcal{P}$ is the powerset functor. This definition leaves equilibria abstract - they may maximise utility from a choice of move or, as in say the El Farrol Bar game, they may reflect other criteria. (The functions E are precisely the multi-valued selection functions [9].) In usual game theory, we have a fixed utility function and a set of equilibria. In our more general setting, we have have equilibria for *arbitrary* utility functions (of the correct type). That is, an equilibrium $y \in Ek$ means that if the utility in the game is given by k, then there is no move preferable to y. And, crucially, the equilibria of the game are not given for a specific utility function, but are given for every utility function! Now, recall the essence of compositionality is to build new games from old in such a way that properties of the former can be derived from those of the later. We now introduce such an operator

Definition 2.3 (Monoidal Product of Simple Games)**.** Let $G_1 \in \mathsf{SP}_{R_1} Y_1$ and $G_2 \in \mathsf{SP}_{R_2} Y_2$ be simple games. Their monoidal product is the simple game $G_1 \otimes G_2 : \mathsf{SP}_{R_1 \times R_2} Y_1 \times Y_2$ with equilibrium function

$$\begin{aligned}(y_1, y_2) \in E_{G_1 \otimes G_2} k \quad \mathsf{iff} \quad & y_1 \in E_1(\pi_1 \circ k(-, y_2)) \\ & \wedge\, y_2 \in E_2(\pi_2 \circ k(y_1, -))\end{aligned}$$

Although the Nash equilibria of a simple 2-player game appeared initially to be a primitive and non-compositional concept, we can now show it arises compositionally. The proof is by unwinding definitions.

Theorem 2.4. *Let $G = (Y_1, Y_2, R_1, R_2, k)$ be a simple 2-player game. Then $(y_1, y_2) \in E_G$ iff $(y_1, y_2) \in E_{G_1 \otimes G_2} k$ where G_1 and G_2 are the simple games $G_1 = (Y_1, R_1, \mathsf{argmax})$ and $G_2 = (Y_2, R_2, \mathsf{argmax})$.*

Note that in the simple games G_1 and G_2, the 'rational behaviour' of the players is to maximise (where 'rational' is taken to mean 'playing according to their equilibrium'). However the rational behaviour of players in $G_1 \otimes G_2$ may not be globally maximising – this is precisely what the Prisoner's Dilemma illustrates. Thus maximising behaviour is not compositional, so the generalisation from argmax to the equilibrium function E is essential. Also notice how in Theorem 2.4 it is essential to not hardwiring a utility function into a simple game but rather define the equilibria for *all* possible utility functions. Finally, notice how the equilibria of $G_1 \otimes G_2$ can be computed compositionally from only the equilibria of G_1 and G_2, even if G_1 and G_2 are highly complex games. That is, one need not delve into the definitions of G_1 and G_2 to see how they are built. In the rest of this paper, we generalise this basic compositional model economic game theory to a more sophisticated compositional model encompassing a much broader class of games.

3. Pregames

Section 2 defined a compositional model of economic game theory. That is, we defined i) the notion of a simple game ii) an operator for building new simple games from old; and iii) the equilibria for compound simple games in terms of the equilibria of their component simple games. However, simple games possess limited structure and hence support limited operators — this is a problem, as more operators enable more compositional reasoning! Therefore, we introduce a more sophisticated compositional model which finds a sweet spot of being expressive enough to model complex and highly irregular games while retaining enough simplicity to continue to deliver conceptual clarity. As motivation, consider me placing a bet with a bookmaker:

- My state informs my actions, e.g. my wealth affects the size of my bet. Secondly, I have different strategies, e.g. high-risk, long-odds bets, or low-risk, short-odds bets. Strategy and state are conceptually different as I can choose my strategy but not my state. Together, a state and a strategy determine the action taken, i.e. which bet I place. The environment, i.e. the bookmaker, returns my winnings.
- The bookmaker also has a state: in this case, the bet placed. They too have a number of strategies, e.g. to refuse the bet, to accept the bet, or to lay the bet. Again, their state and strategy determine their action. And, again, it is the environment that converts their actions into results and hence utility. Finally, the bookmaker will return winnings to the better — we call this *coutility* since it is the source of the better's utility.

The above example motivates our more general definition:

Definition 3.1 (Pregame)**.** Let X, Y, R and S be sets. A *pregame* G over X, Y, R, S consists of the following data:

- A set $\Sigma_{\mathcal{G}}$ of *strategy profiles*
- A *play* function $P_{\mathcal{G}} : \Sigma_{\mathcal{G}} \to (X \to Y)$
- A *coutility* function $C_{\mathcal{G}} : \Sigma_{\mathcal{G}} \to (X \times R \to S)$
- An *equilibrium function* $E_{\mathcal{G}} : X \times (Y \to R) \to \mathcal{P}\Sigma_{\mathcal{G}}$.

For reasons that will become apparent later, we denote such a pregame $\mathcal{G} : (X, S) \to (Y, R)$. We chose the name pregames for the above concept so that we could reserve the term games for the special case of pregames $G : (X, S) \to (1, 1)$, ie pregames for which no further interaction with the environment takes place. All simple games are pregames where $X = S = 1$, $\Sigma = Y$ and P is the identity function. If $\sigma \in E_G(x, k)$, then we imagine there is no better strategy than σ if we are in state x and $k : Y \to R$ computes the utilities arising from each action. We also represent pregames as string diagrams containing ports where interaction with other games takes place.



Example 3.2 (Prisoners Dilemma)**.** Let M be the set of moves in the Prisoners Dilemma game, i.e. $M = \{C, D\}^2$. Define the pregame $\mathsf{PD} : (1, \mathbb{R} \times \mathbb{R}) \to (M, \mathbb{R} \times \mathbb{R})$ with strategies

$\Sigma_{\mathsf{PD}} = M$ and with play function (eliding the unit type) choosing the strategy, i.e.

$$P_{\mathsf{PD}}\ m = m$$

Coutility is $C_{\mathsf{PD}}\ m\ r = r$, while

$$\begin{aligned}(\sigma_1, \sigma_2) \in E_{\mathsf{PD}} k \quad \mathsf{iff} \quad & \sigma_1 \in \mathsf{argmax}\ \pi_1 \circ k(-, \sigma_2) \\ & \wedge\ \sigma_2 \in \mathsf{argmax}\ \pi_2 \circ k(\sigma_1, -)\end{aligned}$$

It is worth remarking that our notion of Prisoner's Dilemma differs from the standard one found in the literature we we leave open the pay-off matrix rather than hardwire one in. We can also define a two round Prisoners Dilemma:

Example 3.3 (2-Round Prisoners Dilemma)**.** Again, let $M = \{C, D\}^2$ be the set of moves in the Prisoners Dilemma game. We define a pregame $\mathsf{2PD} : (1, \mathbb{R} \times \mathbb{R}) \to (M \times M, \mathbb{R} \times \mathbb{R})$ which represents two rounds of the Prisoners Dilemma Game. The strategy set for the game ought to be a move for the first round together with a function which gives a round 2 move for each round 1 moves. That is $\Sigma_{\mathsf{2PD}} = M \times (M \to M)$. The play function is

$$P_{\mathsf{2PD}}\ (\sigma_1, \sigma_2) = (\sigma_1, \sigma_2 \sigma_1)$$

Coutility is similar to that above, while

$$\begin{aligned}(\sigma_1, \sigma_2) \in E_{\mathsf{2PD}}\ k\ \mathsf{iff}\ & \sigma_1 \in E_{\mathsf{PD}}\ k(-, \sigma_2 -) \quad \wedge \\ & \forall \sigma \in M.\ \sigma_2 \sigma \in E_{\mathsf{PD}}\ k(\sigma, -)\end{aligned}$$

Note σ_2 must compute an optimal strategy for all round-1 plays — this accords with the no-incredible-threats principle of subgame perfect equilibria. The above definition of $\mathsf{2PD}$ is non-compositional in that it is not built from smaller games but rather defined from scratch. We now show how we can define operators that build complex pregames from simpler pregames. We shall use these operators to show how $\mathsf{2PD}$ can be built compositionally

4. Operators on Pregames

We introduce operators for building pregames. The first three build atomic pregames while the last two build compound pregames from other pregames:

Definition 4.1 (Players)**.** A player who observes a state of type X, makes a choice of type Y and optimises an outcome of type R is represented by a pregame $\mathcal{P} : (X, 1) \to (Y, R)$, where the set of strategies is $\Sigma_{\mathcal{P}} = X \to Y$, i.e. mappings from states to choices. The play function $P_{\mathcal{P}}(\sigma, x) = \sigma(x)$ applies the strategy to the state, while the coutility function is trivial. In general, the equilibrium function can be varied in different examples to give different goals to the player.

If the player does not make any observation then $X = 1$. In the two cases $X \neq 1$ and $X = 1$, a player is represented by a string diagram as

(triangles traditionally denote components of string diagrams that have connections on only one side). For an example of a classical, utility-maximising player, take $R = \mathbb{R}$ with the equilibrium relation being given by

$$E_{\mathcal{P}}\ (x, k) = \{\sigma : X \to Y \mid \sigma(x) \in \mathsf{argmax}\ k\}$$

If $X = 1$, this simplifies to

$$E_{\mathcal{P}}\ (*, k) = \mathsf{argmax}\ k$$

Definition 4.2 (Computations)**.** A function $f : X \to Y$ defines a two pregames, a 'covariant computation' $f : (X, 1) \to (Y, 1)$ and a 'contravariant computation' $f^* : (1, Y) \to (1, X)$. These are used to represent aspects of a game that are not players (more precisely, that cannot make strategic choices), e.g. utility functions. In both cases we have $\Sigma = 1$, and define E_f and E_{f^*} to always return $\{*\}$. In the covariant case, we set $P_f(*, x) = fx$, while in the contravariant case we set $C_{f^*}(*, *, x) = fx$.

Covariant and contravariant computations are respectively drawn as

An important example of a covariant computation is copying $\Delta_X : (X, 1) \to (X \times X, 1)$ arising from the function $\Delta_X : X \to X \times X$ which copies its argument. Within games, this allows information to be used more than once, for example both being observed by a player, and used as input to a utility function.



Our final basic constructor for pregames is the *Teleological Counit* which mediates between the forward direction of information (chose strategy, play move) and the backward flow of information (a process generates coutility and passes it to a previous process as utility). More concretely, the teleological unit — in combination with a computation — allows us to specify a particular utility function to be used in the pregame. Of course, this curtails the possibility of interacting with other pregames by closing off the ports where that interaction can happen:

Definition 4.3 (Teleological counit)**.** The pregame $\tau_X : (X, X) \to (1, 1)$ is given by $\Sigma_{\tau_X} = 1$, $C_{\tau_X}(*, x, *) = x$ and with E_{τ_X} always returning $\{*\}$.

We graphically represent the teleological counit by a *cup*



After introducing three basic operators for pregames, we now present two compound operators — the monoidal product which generalises the similar operator of Section 2, and sequential composition.

Definition 4.4 (Monoidal Product of Pregames)**.** If $\mathcal{G} : (X_1, R_1) \to (Y_1, S_1)$ and $\mathcal{H} : (X_2, S_2) \to (Y_2, R_2)$ are pregames, their monoidal product

$$\mathcal{G} \otimes \mathcal{H} : (X_1 \times X_2, S_1 \times S_2) \to (Y_1 \times Y_2, R_1 \times R_2)$$

is defined by

- $\Sigma_{\mathcal{G}\otimes\mathcal{H}} = \Sigma_{\mathcal{G}} \times \Sigma_{\mathcal{H}}$
- $P_{\mathcal{G}\otimes\mathcal{H}}(\sigma_1, \sigma_2)(x_1, x_2) = (P_{\mathcal{G}}\ \sigma_1\ x_1, P_{\mathcal{H}}\ \sigma_2\ x_2)$
- $C_{\mathcal{G}\otimes\mathcal{H}}(\sigma_1, \sigma_2)((x_1, x_2), (r_1, r_2)) = (C_{\mathcal{G}}(\sigma_1(x_1, r_1), C_{\mathcal{H}}(\sigma_2)(x_2, r_2))$
- $(\sigma_1, \sigma_2) \in E_{\mathcal{G}\otimes\mathcal{H}}\ ((x_1, x_2),\ k)$ iff $\sigma_1 \in E_{\mathcal{G}}\ (x_1,\ k_1)$ and $\sigma_2 \in E_{\mathcal{H}}\ (x_2,\ k_2)$ where

$$k_1(y_1) = (\pi_1 \circ k)(y_1, P_{\mathcal{H}}(\sigma_2)(x_2))$$
$$k_2(y_2) = (\pi_2 \circ k)(P_{\mathcal{G}}(\sigma_1)(x_1), y_2)$$

Definition 4.5 (Sequential Composition)**.** Let $\mathcal{G} : (X, T) \to (Y, S)$ and $\mathcal{H} : (Y, S) \to (Z, R)$ be pregames. The composition $\mathcal{H} \circ \mathcal{G} : (X, T) \to (Z, R)$ is defined by

- $\Sigma_{\mathcal{H}\circ\mathcal{G}} = \Sigma_{\mathcal{G}} \times \Sigma_{\mathcal{H}}$
- $P_{\mathcal{H}\circ\mathcal{G}}(\sigma_1, \sigma_2) = P_{\mathcal{H}}(\sigma_2) \circ P_{\mathcal{G}}(\sigma_1)$
- $C_{\mathcal{H}\circ\mathcal{G}}(\sigma_1, \sigma_2)(x, r) = C_{\mathcal{G}}(\sigma_1)(x, C_{\mathcal{H}}(\sigma_2)(P_{\mathcal{G}}(\sigma_1)(x), r)))$
- $(\sigma_1, \sigma_2) \in E_{\mathcal{H}\circ\mathcal{G}}(x, k)$ iff
  (1) $\sigma_1 \in E_{\mathcal{G}}(x, k')$, where $k'(y) = C_{\mathcal{H}}(\sigma_2)(y, k(P_{\mathcal{H}}(\sigma_2)(y)))$, and
  (2) $\sigma_2 \in E_{\mathcal{H}}(P_{\mathcal{G}}(\sigma_1', x), k)$ for all $\sigma_1' \in \Sigma_{\mathcal{G}}$

The definition of sequential composition highlights two fundamental points

- We promised to explain why we write $\mathcal{G} : (X, S) \to (Y, R)$ if $\mathcal{G}$ has state X, moves Y, utility R and coutility S. When composing functions $f : A \to B$ and $g : B \to C$, the codomain of f must be the same as the domain of g because we feed the output of f into the input of g. Similarly, in $\mathcal{H} \circ \mathcal{G}$, we feed the output move $\mathcal{G}$ into the state of $\mathcal{H}$, and the coutility of $\mathcal{H}$ into the utility of $\mathcal{G}$. Thus i) the type of moves of $\mathcal{G}$ and the states of $\mathcal{H}$ must be the same; and ii) the type of utility of $\mathcal{G}$ and the coutility of $\mathcal{H}$ must also be

the same. Thus, the domain of a game contains its state and coutility, while its codomain contains its moves and utility.
- To treat the composition $\mathcal{H} \circ \mathcal{G}$ correctly - as in the example of placing a bet with a bookmaker - $\mathcal{G}$ can only get its utility once H has its own utility. Technically, within the definition of equilibrium for a composed system, $\mathcal{G}$ must be in equilibrium for a utility function $Y \to R$ which depends upon what $\mathcal{H}$ feeds back to $\mathcal{G}$. This ability of the future system $\mathcal{H}$ to feed utility into a previous system $\mathcal{G}$ is exactly why we introduced the concept of coutility. Formally, the definition of equilibrium in the composition $E_{\mathcal{H}\circ\mathcal{G}}$ depends upon the coutility $C_{\mathcal{H}}$.

Example 4.6. A two-player context-dependent game is defined in [9] to consist of the following data:
- Sets X, Y of choices for each player, and R of outcomes
- Multivalued selection functions $\varepsilon : (X \to R) \to \mathcal{P}X$, $\delta : (Y \to R) \to \mathcal{P}Y$
- An outcome function $q : X \times Y \to R$
- A strategy profile is simply a pair $(\sigma_1, \sigma_2) : X \times Y$. A strategy profile is called a *selection equilibrium* if

$$\sigma_1 \in \varepsilon(q(-, \sigma_2))$$
$$\sigma_2 \in \delta(q(\sigma_1, -))$$

Context dependent games generalise the simple games of Def 2.2 in that the selection functions need not be argmax. For example, in [9] it is shown that selection functions returning a set of fixpoints or non-fixpoints gives an elegant model of coordination or differentiation goals of players, respectively. The selection equilibria of context dependent games can be characterised compositionally.

Theorem 4.7. *The selection equilibria of this game are precisely the equilibria of the string diagram*



where q is the lifting of the outcome function as a covariant computation, and $\mathcal{P}_\varepsilon$ and $\mathcal{P}_\delta$ are players, whose equilibrium functions are respectively

$$E_{\mathcal{P}_\varepsilon}(*, k) = \varepsilon k$$
$$E_{\mathcal{P}_\delta}(*, k) = \delta k$$

Proof. The above string diagram is clearly definable within our grammar of pregames, for example, by the expression

$$\mathcal{G} = (\tau_R \otimes \tau_R) \circ (\mathrm{id}_R^* \otimes (\Delta_R \circ q) \otimes \mathrm{id}_R^*) \circ (\mathcal{P}_\varepsilon \otimes \mathcal{P}_\delta) : (1, 1) \to (1, 1)$$

Unwinding the definitions, we have

$$\begin{aligned} & (\sigma_1, \sigma_2) \in E_{\mathcal{G}}(*,*) \\ \iff & (\sigma_1, \sigma_2) \in E_{\mathcal{P}_\varepsilon \otimes \mathcal{P}_\delta}(*, k) \wedge * \in E_{(\tau_R \otimes \tau_R) \circ (\mathrm{id}_R^* \otimes (\Delta_R \circ q) \otimes \mathrm{id}_R^*)}(*,*) \end{aligned}$$

The second term of this is found to be vacuously true, and so it is equivalent to

$$\begin{aligned} & \iff (\sigma_1, \sigma_2) \in E_{\mathcal{P}_\varepsilon \otimes \mathcal{P}_\delta}(*, k) \\ & \iff \sigma_1 \in E_{\mathcal{P}_\varepsilon}(*, k_1) \wedge \sigma_2 \in E_{\mathcal{P}_\delta}(*, k_2) \\ & \iff \sigma_1 \in \varepsilon k_1 \wedge \sigma_2 \in \delta k_2 \end{aligned}$$

It is straightforward but tedious to verify that, according to the above definitions, the continuations k, k_1 and k_2 are given as follows:

$$\begin{aligned} k(x,y) &= C_{(\tau_R \otimes \tau_R) \circ (\mathrm{id}_R^* \otimes (\Delta_R \circ q) \otimes \mathrm{id}_R^* \otimes \mathrm{id}_R^*)}(*)((x,y),*) \\ &= (q(x,y), q(x,y)) \\ k_1(x) &= (\pi_1 \circ k)(x, P_{\mathcal{P}_\delta}(\sigma_2, *)) = (\pi_1 \circ k)(x, \sigma_2) = q(x, \sigma_2) \\ k_2(y) &= (\pi_2 \circ k)(P_{\mathcal{P}_\varepsilon}(\sigma_1, *), y) = (\pi_2 \circ k)(\sigma_1, y) = q(\sigma_1, y) \end{aligned}$$

Thus, we have proved that $E_{\mathcal{G}}(*,*)$ is exactly the set of (σ_1, σ_2) that satisfy the conditions of a selection equilibrium. □

Context dependent games and selection equilibria include ordinary normal form games and Nash equilibria as a special case, by taking $R = \mathbb{R}^2$, and the selection function ε and δ to be the argmax operators for the first and second coordinates:

$$\begin{aligned} \varepsilon(k) &= \{x \in X \mid (\pi_1 \circ k)(x) \geq (\pi_1 \circ k)(x') \text{ for all } x' \in X\} \\ \delta(k) &= \{y \in Y \mid (\pi_2 \circ k)(y) \geq (\pi_2 \circ k)(y') \text{ for all } y' \in Y\} \end{aligned}$$

A second class of games that are subsumed by our grammar are generalised sequential games, which subsume classical games of perfect information.

Definition 4.8. A two-player sequential game is defined in [7] to consist of the following data:

- Sets X, Y of choices for each player, and R of outcomes
- Multivalued quantifiers $\varphi : (X \to R) \to \mathcal{P}R$, $\psi : (Y \to R) \to \mathcal{P}R$
- An outcome function $q : X \times Y \to R$
- A strategy profile for this game consists of a move $\sigma_1 : X$ for the first player and a contingent strategy $\sigma_2 : X \to Y$ for the second player. A strategy profile is called *optimal* if

$$\begin{aligned} q(\sigma_1, \sigma_2(\sigma_1)) &\in \varphi \lambda x. q(x, \sigma_2(x)) \\ q(x, \sigma_2(x)) &\in \psi \lambda y. q(x,y) \quad \text{for all } x \in X \end{aligned}$$

Note that the difference between selection functions and quantifiers is relatively unimportant: we could equally well define simultaneous games using quantifiers, and sequential games using selection functions.

Theorem 4.9. *Every optimal strategy profile for this game is an equilibrium of the string diagram*



where P_ε and P_δ are players with equilibrium functions

$$\sigma_1 \in E_{P_\varphi}(*, k_1) \iff k_1(\sigma_1) \in \varphi(k_1)$$
$$\sigma_2 \in E_{P_\psi}(x, k_2) \iff k_2(\sigma_2(x)) \in \psi(k_2)$$

Proof. Again, this string diagram falls within our grammar being given algebraically by

$$(\tau_R \otimes \tau_R) \circ (\mathrm{id}_R^* \otimes (((\Delta_R \circ q) \otimes \mathrm{id}_R^*) \circ (\mathrm{id}_X \otimes \mathcal{P}_\psi) \circ \Delta_X)) \circ \mathcal{P}_\varphi : (1,1) \to (1,1)$$

Unwinding the definition, we have

$$\begin{aligned} & (\sigma_1, \sigma_2) \in E_{(\tau_R \otimes \tau_R) \circ (\mathrm{id}_R^* \otimes (((\Delta_R \circ q) \otimes \mathrm{id}_R^*) \circ (\mathrm{id}_X \otimes \mathcal{P}_\psi) \circ \Delta_X)) \circ \mathcal{P}_\varphi}(*, *) \\ \iff & (\sigma_1, \sigma_2) \in E_{(\mathrm{id}_R^* \otimes (((\Delta_R \circ q) \otimes \mathrm{id}_R^*) \circ (\mathrm{id}_X \otimes \mathcal{P}_\psi) \circ \Delta_X)) \circ \mathcal{P}_\varphi}(*, \mathrm{id}_{R \times R}) \\ \iff & \sigma_1 \in E_{\mathcal{P}_\varphi}(*, k_1) \ \ \wedge \ \ \forall \sigma_1' \in X.\ \ \sigma_2 \in E_{\mathcal{G}}(P_{\mathcal{P}_\varphi}(\sigma_1')(*), \mathrm{id}_{R \times R}) \end{aligned}$$

where

$$\mathcal{G} = \mathrm{id}_R^* \otimes (((\Delta_R \circ q) \otimes \mathrm{id}_R^*) \circ (\mathrm{id}_X \otimes \mathcal{P}_\psi) \circ \Delta_X) : (X, R) \to (R \times R, R \times R)$$

and

$$k_1(x) = C_{\mathcal{G}}(\sigma_2)(x, \mathrm{id}_{R \times R}(P_{\mathcal{G}}(\sigma_2)(x))) = q(x, \sigma_2(x))$$

The first condition is therefore equivalent to

$$q(\sigma_1, \sigma_2(\sigma_1)) \in \varphi(\lambda x. q(x, \sigma_2(x)))$$

which is the first part of the definition of an optimal strategy.

Continuing, the second condition for a particular $\sigma_1' \in X$ is

$$\begin{aligned}
\sigma_2 \in E_{\mathcal{G}}(P_{\mathcal{P}_\varepsilon}(\sigma_1')(*), \mathrm{id}_{R\times R}) &\iff \sigma_2 \in E_{\mathcal{G}}(\sigma_1', \mathrm{id}_{R\times R}) \\
&\iff \sigma_2 \in E_{((\Delta_R \circ q)\otimes \mathrm{id}_R^*)\circ(\mathrm{id}_X \otimes \mathcal{P}_\psi)\circ\Delta_X}(\sigma_1', \pi_2) \\
&\iff \sigma_2 \in E_{((\Delta_R \circ q)\otimes \mathrm{id}_R^*)\circ(\mathrm{id}_X \otimes \mathcal{P}_\psi)}((\sigma_1', \sigma_1'), \pi_2) \\
&\iff \sigma_2 \in E_{\mathrm{id}_X \otimes \mathcal{P}_\psi}((\sigma_1', \sigma_1'), k_2) \\
&\iff \sigma_2 \in E_{\mathcal{P}_\psi}(\sigma_1', k_3)
\end{aligned}$$

where

$$\begin{aligned}
k_2(x,y) &= C_{(\Delta_R \circ q)\otimes \mathrm{id}_R^*}(*)((x,y), (\pi_2 \circ P_{(\Delta_R \circ q)\otimes \mathrm{id}_R^*}(*))(x,y))) \\
&= q(x,y) \\
k_3(y) &= k_2(\sigma_1', y) = q(\sigma_1', y)
\end{aligned}$$

Thus, the second condition is that for all $\sigma_1' \in X$,

$$k_3(\sigma_2(\sigma_1')) \in \psi(k_3)$$

which is

$$q(\sigma_1', \sigma_2(\sigma_1')) \in \psi(\lambda y. q(\sigma_1', y))$$

which is the second part of the definition of an optimal strategy. □

Finally we can see 2PD as being built compositionally from PD. Note that this is not as one might think done by simply gluing tow copies of PD together as we must condition the second round of PD on the move arising in he first round of PD. First, given a pregame $G : (X, S) \to (Y, R)$ with strategies Σ and a set A, define the pregame $A \to G : (A\times X, S) \to (Y, R)$ to have strategies given by $A \to \Sigma$ and the obvious play, coutility and equilibria. Then 2PD arises by composing PD with $M \to$ PD and copying the move from the first round to the move of the second round. Formally, 2PD arises as

$$((M \to \mathsf{PD}) \otimes id) \circ (id^* \otimes \Delta_M) \circ \mathsf{PD}$$

where, recall, $\Delta : M \to M \times M$.

5. The Monoidal Category of Pregames

The mathematical treatment of pregames above delivers a bottom-up collection of operators for defining complex pregames from simple pregames and computing the equilibria of those complex games from their more simple components. Furthermore, pregames naturally posses a graphical form as a string diagram showing the flow of information around the game. This makes pregames easier to visualise and therefore comprehend. However, this mathematical treatment is so far insufficient for a number of reasons:

- Reasoning about string diagrams needs us to formally define diagrams — this can be a very intricate and messy process. Category theory offers a treatment of string diagrams known as *monoidal category theory* which has been successfully applied to a number of areas, eg network theory, quantum physics, concurrency theory.
- We want our operators to capture fundamental structure. Category theory is used to find fundamental structures across mathematics and computer science, eg one such is the monoidal product which arises in both simple games and pregames.

- To reason effectively, we need to know when two different pregames are the same, eg the associativity of composition or the distributivity of contra variant computations over composition: $(g \circ f)^* = f^* \circ g^*$. Category theory gives canonical laws one can expect to hold between the canonical operators that arise in category theory.
- Category theory provides universal properties for structuring and reasoning. A key example are final coalgebras which we use to model infinite iteration of pregames.

For these reasons (and for others more related to future work), we introduce a categorical model of pregames. The first guess is to take as objects pairs of sets and define the morphisms from (X, S) to (Y, R) to be the pregames $(X, S) \to (Y, R)$. However this doesn't quite work, e.g. composition would fail to be associative. The problem is that games which have isomorphic sets of strategies ought really to be the same and hence we quotient pregames by such isomorphisms. This is in tune with general categorical principles where one allows objects to be isomorphic but asks morphisms to either be equal or not. A similar approach underlies the construction of free cartesian closed categories where one quotients well typed λ-calculus terms by $\beta\eta$-equality.

Theorem 5.1. *There is a category* Pregame *whose objects are pairs of sets (X, S). Morphisms from (X, S) to (Y, R) are pregames $(X, S) \to (Y, R)$ quotiented by the relation which identifies two pregames if their sets of strategy profiles are isomorphic and their play, coutility and equilibrium functions are the same under this isomorphism.*

Proof. To ensure readability, our proof treats pregames and we omit the entirely trivial checks that our arguments preserve the equivalence relation on pregames. On the other hand, since morphisms/pregames contain a lot of data, we give the rest of the proof in some detail. The identity on the object (X, S) is the pregame $\mathrm{id}_X \otimes \mathrm{id}_S^*$. More concretely, this is the pregame

- $\Sigma_{\mathrm{id}_{(X,S)}} = 1$
- $P_{\mathrm{id}_{(X,S)}}(*) = \mathrm{id}_X$
- $C_{\mathrm{id}_{(X,S)}}(*)(x, r) = r$
- $* \in E_{\mathrm{id}_{(X,S)}}(x, k)$ for every x and k

Left identity. Let $\mathcal{G} : (X, S) \to (Y, R)$. We prove that $\mathrm{id}_{(Y,R)} \circ \mathcal{G} = \mathcal{G}$. Then

- $\Sigma_{\mathrm{id}_{(Y,R)} \circ \mathcal{G}} = \Sigma_{\mathrm{id}_{(Y,R)}} \times \Sigma_{\mathcal{G}} = 1 \times \Sigma_{\mathcal{G}} = \Sigma_{\mathcal{G}}$
- $P_{\mathrm{id}_{(Y,R)} \circ \mathcal{G}}(\sigma) = P_{\mathrm{id}_{(Y,R)}}(*) \circ P_{\mathcal{G}}(\sigma) = \mathrm{id}_Y \circ P_{\mathcal{G}}(\sigma) = P_{\mathcal{G}}(\sigma)$
- $C_{\mathrm{id}_{(Y,R)} \circ \mathcal{G}}(\sigma)(x, r) = C_{\mathcal{G}}(\sigma)(x, C_{\mathrm{id}_{(Y,R)}}(*)(P_{\mathcal{G}}(\sigma)(x), r)) = C_{\mathcal{G}}(\sigma)(x, r)$
- $\sigma \in E_{\mathrm{id}_{(Y,R)} \circ \mathcal{G}}(x, k) \iff \sigma \in E_{\mathcal{G}}(x, k') \wedge * \in E_{\mathrm{id}_{(Y,R)}}(P_{\mathcal{G}}(\sigma)(x), k) \iff \sigma \in E_{\mathcal{G}}(x, k')$ where
$$k'(y) = C_{\mathrm{id}_{(Y,R)}}(*)(y, k(P_{\mathrm{id}_{(Y,R)}}(*)(y))) = k(P_{\mathrm{id}_{(Y,R)}}(*)(y)) = k(y)$$

Right identity. We prove that $\mathcal{G} \circ \mathrm{id}_{X,S} = \mathcal{G}$. We have

- $\Sigma_{\mathcal{G} \circ \mathrm{id}_{(X,S)}} = \Sigma_{\mathcal{G}} \times \Sigma_{\mathrm{id}_{(X,S)}} = \Sigma_{\mathcal{G}} \times 1 = \Sigma_{\mathcal{G}}$
- $P_{\mathcal{G} \circ \mathrm{id}_{(X,S)}}(\sigma) = P_{\mathcal{G}}(\sigma) \circ P_{\mathrm{id}_{(X,S)}}(*) = P_{\mathcal{G}}(\sigma) \circ \mathrm{id}_X = P_{\mathcal{G}}(\sigma)$
- $C_{\mathcal{G} \circ \mathrm{id}_{(X,S)}}(\sigma)(x, r) = C_{\mathrm{id}_{(X,S)}}(*)(x, C_{\mathcal{G}}(\sigma)(P_{\mathrm{id}_{(X,S)}}(*)(x), r)) =$
  $C_{\mathcal{G}}(\sigma)(P_{\mathrm{id}_{(X,S)}}(*)(x), r) = C_{\mathcal{G}}(\sigma)(x, r)$
- $\sigma \in E_{\mathcal{G} \circ \mathrm{id}_{(X,S)}}(x, k) \iff * \in E_{\mathrm{id}_{(X,S)}}(x, k') \wedge \sigma \in E_{\mathcal{G}}(P_{\mathrm{id}_{(X,S)}}(*)(x), k)$
  $\iff \sigma \in E_{\mathcal{G}}(x, k)$

Associativity. Let $\mathcal{G} : (X, U) \to (Y, T)$, $\mathcal{H} : (Y, T) \to (Z, S)$ and $\mathcal{I} : (Z, S) \to (W, R)$. We have

$$\Sigma_{(\mathcal{I}\circ\mathcal{H})\circ\mathcal{G}} = \Sigma_{\mathcal{G}} \times \Sigma_{\mathcal{I}\circ\mathcal{H}} = \Sigma_{\mathcal{G}} \times \Sigma_{\mathcal{H}} \times \Sigma_{\mathcal{I}} = \Sigma_{\mathcal{H}\circ\mathcal{G}} \times \Sigma_{\mathcal{I}} = \Sigma_{\mathcal{I}\circ(\mathcal{H}\circ\mathcal{G})}$$

For the play function we have

$$\begin{aligned} P_{(\mathcal{I}\circ\mathcal{H})\circ\mathcal{G}}(\sigma_1, \sigma_2, \sigma_3) &= P_{\mathcal{I}\circ\mathcal{H}}(\sigma_2, \sigma_3) \circ \mathbb{P}_{\mathcal{G}}(\sigma_1) \\ &= P_{\mathcal{I}}(\sigma_3) \circ P_{\mathcal{H}}(\sigma_2) \circ P_{\mathcal{G}}(\sigma_1) \\ &= P_{\mathcal{I}}(\sigma_3) \circ P_{\mathcal{H}\circ\mathcal{G}}(\sigma_1, \sigma_2) \\ &= P_{\mathcal{I}\circ(\mathcal{H}\circ\mathcal{G})}(\sigma_1, \sigma_2, \sigma_3) \end{aligned}$$

and for the coutility function have

$$\begin{aligned} & C_{(\mathcal{I}\circ\mathcal{H})\circ\mathcal{G}}(\sigma_1, \sigma_2, \sigma_3)(x, r) \\ = {} & C_{\mathcal{G}}(\sigma_1)(x, C_{\mathcal{I}\circ\mathcal{H}}(\sigma_2, \sigma_3)(P_{\mathcal{G}}(\sigma_1)(x), r)) \\ = {} & C_{\mathcal{G}}(\sigma_1)(x, C_{\mathcal{H}}(\sigma_2)(P_{\mathcal{G}}(\sigma_1)(x), C_{\mathcal{I}}(\sigma_3)(P_{\mathcal{H}}(\sigma_2)(P_{\mathcal{G}}(\sigma_1)(x), r)))) \\ = {} & C_{\mathcal{G}}(\sigma_1)(x, C_{\mathcal{H}}(\sigma_2)(P_{\mathcal{G}}(\sigma_1)(x), C_{\mathcal{I}}(\sigma_3)(P_{\mathcal{H}\circ\mathcal{G}}(\sigma_1, \sigma_2)(x), r))) \\ = {} & C_{\mathcal{H}\circ\mathcal{G}}(\sigma_1, \sigma_2)(x, C_{\mathcal{I}}(\sigma_3)(P_{\mathcal{H}\circ\mathcal{G}}(\sigma_1, \sigma_2)(x), r)) \\ = {} & C_{\mathcal{I}\circ(\mathcal{H}\circ\mathcal{G})}(\sigma_1, \sigma_2, \sigma_3)(x, r) \end{aligned}$$

For the equilibrium condition we have

$$\begin{aligned} & (\sigma_1, \sigma_2, \sigma_3) \in E_{(\mathcal{I}\circ\mathcal{H})\circ\mathcal{G}}(x, k_3) \\ \iff {} & \sigma_1 \in E_{\mathcal{G}}(x, k_1) \wedge \forall \sigma_1' \in \Sigma_{\mathcal{G}}.(\sigma_2, \sigma_3) \in E_{\mathcal{I}\circ\mathcal{H}}(P_{\mathcal{G}}(\sigma_1')(x), k_3) \\ \iff {} & \sigma_1 \in E_{\mathcal{G}}(x, k_1) \wedge \forall \sigma_1' \in \Sigma_{\mathcal{G}}.[\sigma_2 \in E_{\mathcal{H}}(P_{\mathcal{G}}(\sigma_1')(x), k_2) \\ & \wedge \forall \sigma_2' \in \Sigma_{\mathcal{H}}.\sigma_3 \in E_{\mathcal{I}}(P_{\mathcal{H}}(\sigma_2')(P_{\mathcal{G}}(\sigma_1')(x)), k_3)] \\ \iff {} & \sigma_1 \in E_{\mathcal{G}}(x, k_1) \wedge [\forall \sigma_1' \in \Sigma_{\mathcal{G}}.\sigma_2 \in E_{\mathcal{H}}(P_{\mathcal{G}}(\sigma_1')(x), k_2)] \\ & \wedge \forall (\sigma_1', \sigma_2') \in \Sigma_{\mathcal{H}\circ\mathcal{G}}.\sigma_3 \in E_{\mathcal{I}}(P_{\mathcal{H}\circ\mathcal{G}}(\sigma_1', \sigma_2')(x), k_3) \\ \iff {} & (\sigma_1, \sigma_2) \in E_{\mathcal{H}\circ\mathcal{G}}(x, k_2) \\ & \wedge \forall (\sigma_1', \sigma_2') \in \Sigma_{\mathcal{H}\circ\mathcal{G}}.\sigma_3 \in E_{\mathcal{I}}(P_{\mathcal{H}\circ\mathcal{G}}(\sigma_1', \sigma_2')(x), k_3) \\ \iff {} & (\sigma_1, \sigma_2, \sigma_3) \in E_{\mathcal{I}\circ(\mathcal{H}\circ\mathcal{G})}(x, k_3) \end{aligned}$$

where

$$\begin{aligned} k_1(y) &= C_{\mathcal{I}\circ\mathcal{H}}(\sigma_2, \sigma_3)(y, k_3(P_{\mathcal{I}\circ\mathcal{H}}(\sigma_2, \sigma_3)(x))) \\ &= C_{\mathcal{I}\circ\mathcal{H}}(\sigma_2, \sigma_3)(y, k_3(P_{\mathcal{I}}(\sigma_3)(P_{\mathcal{H}}(\sigma_2)(x)))) \\ &= C_{\mathcal{H}}(\sigma_2)(y, C_{\mathcal{I}}(\sigma_3)(P_{\mathcal{H}}(\sigma_2)(y), k_3(P_{\mathcal{I}}(\sigma_3)(P_{\mathcal{H}}(\sigma_2)(x))))) \\ &= C_{\mathcal{H}}(\sigma_2)(y, k_2(P_{\mathcal{H}}(\sigma_2)(x))) \\ k_2(z) &= C_{\mathcal{I}}(\sigma_3)(z, k_3(P_{\mathcal{I}}(\sigma_3)(z))) \end{aligned}$$

This completes the proof. □

Theorem 5.1 demonstrates how our composition operator is a fundamental algebraic operation. The same is true of our operation for putting games in parallel.

Theorem 5.2. *The category* **Pregame** *is symmetric monoidal.*

Proof. The monoidal product acts on objects (X, R) and (X', R') by taking their product componentwise giving $(X \times X', R \times R')$. The action of the monoidal product on morphisms is given by parallel composition. The unit of this monoidal structure is $(1, 1)$, while the symmetry is inherited from that of the product on the category of sets. As the proof that this data does indeed form a symmetric monoidal category is straightforward, containing none of the detail of Theorem 5.1, we leave the proof as an exercise. □

One final piece of categorical structure bears a striking resemblance to graphical reasoning in a compact closed category:

Theorem 5.3. *If $f : X \to Y$ is a computation then $\tau_Y \circ (f \otimes \mathrm{id}_Y^*) = \tau_X \circ (\mathrm{id}_X \otimes f^*)$. In diagrams,*



Proof. Firstly note that we have

$$P_{f \otimes \mathrm{id}_Y^*}(*)(x) = f(x) \qquad C_{f \otimes \mathrm{id}_Y^*}(*)(x, y) = y$$
$$P_{\mathrm{id}_X \otimes f^*}(*)(x) = x \qquad C_{\mathrm{id}_X \otimes f^*}(*)(x, x') = f(x')$$

We have $\Sigma_{\tau_Y \circ (f \otimes \mathrm{id}_Y^*)} \cong 1 \cong \Sigma_{\tau_X \circ (\mathrm{id}_X \otimes f^*)}$. Both play functions are the unique function $X \to 1$. For the coutility function,

$$C_{\tau_Y \circ (f \otimes \mathrm{id}_Y^*)}(*)(x, *) = f(x) = C_{\tau_X \circ (\mathrm{id}_X \otimes f^*)}(*)(x, *)$$

Finally for both cases we have $* \in E(x, *)$ for all x. □

6. Conclusions and Future Research

We finish the paper with a discussion about what we have achieved and what we intend to do in the future. This paper's most significant contribution is a new way to think about economic games using the idea of *compositionality* from computer science. This involved the development of a new model of economic games which had several original features: i) games do not come with a specific utility function; ii) games can reason about how current actions effect future utility via the new concept of *coutility*; iii) games posses a clear visual interface as string diagrams; and iv) have a fully rigorous treatment within monoidal category theory. The compositionality of this model is demonstrated by the operators that can be defined on arbitrary games and used to build, reason about and even implement larger games from smaller games. Before discussing future work, we analyse decisions implicitly taken is the design of our model and their validity.

- To facilitate accessibility of our paper we chose to work internally to the category of sets as opposed to internally to an ambient symmetric monoidal category. The choice between clarity and mathematical sophistication is important - and we believe we have it right - but is not inherently a scientific one but rather a subjective matter of presentation.
- We made pregames the morphisms of a category to highlight the inputs and outputs of games which was a prerequisite to formalising the idea of *coutility*. This also enabled a string diagram interpretation of pregames thereby giving rigour to our pregame constructors and relating them to constructions on string diagrams. However, ought we to have made pregames the objects of a category? Certainly making pregames objects would have enabled universal properties to have been developed for pregame operators. Actually, both perspectives are valid. Just as the category of sets, relations and relation preserving morphisms is naturally a 2-category, so the category of pregames is naturally a 2-category. For this paper, the 2-dimensional structure was not relevant so we focussed on the 1-dimensional structure of pregames. As commented below, in future work we will need to turn to the 2-dimensional structure of pregames.

Future Work: The potential applications of a compositional, graphical game theory are numerous, especially in economics, and this paper also raises some interesting theoretical questions. We conclude by broadly giving some future research directions and questions:

- Possibly the most important theoretical concepts missing from this paper are incomplete information games, which are ubiquitous in economic applications. This requires the 2-dimensional structure of pre-games alluded to above. Examples of such games show that delicate choices need to be made about whether data should be treated covariantly, contravariantly or relationally. Nevertheless, we have initial thoughts and hope to progress them soon. The 2-dimensional structure will also allow us to tackle infinitely repeated games using final coalgebras.
- Another important aspect of game theory that cannot be modelled by the operators above is the ability of a pregame to depend on a previous move. Influence diagrams also suffer from this problem. For example in a market entry game a firm decides whether to enter a new market, and then subsequently a particular game is played only if the market was entered. This is also commonly used with the 'moves of nature' in the standard approach to incomplete information. Our approach will be to use dependent types and fibred monoidal categories to underpin dependently typed string diagrams.
- Computer support is vital. The first author has developed a Haskell implementation, but it is extremely awkward because the Haskell type system does not unify types like $X \sim 1 \times X$ and $X \sim 1 \to X$, and so the user must manually track these isomorphisms. As an intermediate step, a code generator for a domain specific language similar to Haskell's arrows [14] would be useful. Unfortunately, for technical reasons it does not seem to be possible to use GHC's built-in arrow preprocessor. Ultimately a graphical interface would be invaluable for these ideas to become accessible to working economists.
- As a by-product of obtaining a compositional theory, we have the ability to model preferences of agents which are extremely different to utility maximisation or preference relations. This extends a line of work begun in [9], which uses fixpoint selection functions to model coordination and differentiation. Obvious next steps include modelling bounded rationality [17] and social concerns.
- As described above, a potentially very powerful dimension is to vary the underlying category from Sets to another category. The use of ordinary (possibilistic) nondeterminism in game theory is explored in [15, 4, 8] and [10, chapter 9], and work in progress by the

author suggests that the order structure on possibilistic strategies is important. We also have experimental evidence that correlated equilibria [2] appear as a special case by using a commutative monad transformer stack in which a reader monad gives players read-only access to a shared randomising device. This is strong evidence that side effects in the sense of programming languages can also be a unifying idea in game theory.

- Using noncommutative side-effects is potentially even more rewarding. In this case, the category of pregames may be premonoidal. A major aim is to use strategies with mutable states to model learning, and individual rationality relations to specify that a strategy can be subjectively rational with respect to the current epistemic state, for example using methods of epistemic game theory [16].

References

[1] Samson Abramsky and Viktor Winschel. Coalgebraic analysis of subgame-perfect equilibria in infinite games without discounting. Archiv: https://arxiv.org/abs/1210.4537, 2013.

[2] Robert Aumann. Subjectivity and correlation in randomized strategies. *Journal of mathematical economics*, 1:67–96, 1974.

[3] John Baez and Mike Stay. Physics, topology, logic and computation: a Rosetta stone. In Bob Coecke, editor, *New structures for physics*, pages 95–172. Springer, 2010.

[4] Achim Blumensath and Viktor Winschel. A coalgebraic framework for games in economics. Working paper, 2013.

[5] Filippo Bonchi, Pawel Sobocinski, and Fabio Zanasi. Interacting bialgebras are Frobenius. In *Proceedings of FoSSaCS 2014*, volume 8412, pages 351–365. Springer, 2014.

[6] Bob Coecke, Edward Grefenstette, and Mehrnoosh Sadrzadeh. Lambek vs. Lambek: functorial vector space semantics and string diagrams for Lambek calculus. *Annals of pure and applied logic*, 2013.

[7] Martin Escardó and Paulo Oliva. Sequential games and optimal strategies. *Proceedings of the Royal Society A*, 467:1519–1545, 2011.

[8] Jules Hedges. Monad transformers for backtracking search. In *Proceedings of MSFP'14*, EPTCS, pages 31–50, 2014.

[9] Jules Hedges, Paulo Oliva, Evguenia Sprits, Viktor Winschel, and Philipp Zahn. Higher-order game theory. Submitted, 2015.

[10] Steven M. LaValle. *Planning algorithms*. Cambridge University Press, 2006.

[11] Pierre Lescanne. Rationality and escalation in infinite extensive games. Archiv: https://arxiv.org/abs/1112.1185, 2012.

[12] S MacLane. *Categories for the Working Mathematician*. Springer, 1971.

[13] Paulo Oliva and Thomas Powell. A constructive interpretation of Ramsey's theorem via the product of selecion functions. *Mathematical structures in computer science*, 2014.

[14] Ross Paterson. A new notation for arrows. In *Proceedings of the sixth ACM SIGPLAN international conference on functional programming*, pages 229–240. ACM, 2001.

[15] Dusko Pavlovic. A semantical approach to equilibria and rationality. In *Algebra and coalgebra in computer science*, volume 5728 of *Lectures notes in computer science*, pages 317–334. Springer, 2009.

[16] Andrés Perea. *Epistemic game theory*. Cambridge University Press, 2012.

[17] Ariel Rubinstein. *Modelling bounded rationality*. MIT Press, 1998.

[18] Peter Selinger. A survey of graphical languages for monoidal categories. In Bob Coecke, editor, *New structures for physics*, pages 289–355. Springer, 2011.

This work is licensed under the Creative Commons Attribution-NoDerivs License. To view a copy of this license, visit http://creativecommons.org/licenses/by-nd/2.0/ or send a letter to Creative Commons, 171 Second St, Suite 300, San Francisco, CA 94105, USA, or Eisenacher Strasse 2, 10777 Berlin, Germany